

## **Office of Naval Research Graduate Traineeship Award in Ocean Acoustics for Ankita Deepak Jain**

PI: Prof. Nicholas C. Makris  
Massachusetts Institute of Technology  
77 Massachusetts Avenue, Room 5-212  
Cambridge, MA 02139  
phone: (617) 258-6104 fax: (617) 253-2350 email: [makris@mit.edu](mailto:makris@mit.edu)

Grant Number: N00014-09-1-0679

### **OBJECTIVES**

A primary goal of this research is to model biological clutter in the continental shelf environments of the ocean. We aim to do this by modeling scattered returns from fish shoals in continental shelf environments using a full field matched filter approach as well as its single frequency approximation. This will help in characterizing clutter and will help distinguish scattered fields of moving targets from stationary background reverberation and submerged targets in sonar data.

We aim to develop a unified theory and model for ocean reverberation dependent on seafloor parameters such as density, compressibility and coherence volume. We will use a full field matched filter approach and its time harmonic approximation to model reverberation from volume inhomogeneities. A Monte-Carlo approach based on the parabolic equation will be applied to model acoustic wave propagation in a fluctuating ocean waveguide. We aim at using the model to invert for these parameters and obtain accurate estimates by calibrating modeled returns with data collected during past Ocean Acoustic Waveguide Remote Sensing (OAWRS) experiments in 2003 and 2006 in the New Jersey continental shelf and the Gulf of Maine, respectively.

The final goal of this research is to test the hypothesis that inexpensive underwater acoustic measurements can be used to determine the wind speed and classify the destructive power of a hurricane with greater accuracy than standard satellite remote sensing techniques and with at least the same accuracy as hurricane hunting aircraft.

### **APPROACH**

During the past OAWRS experiments in 2003 and 2006, we demonstrated that fish schools are the dominant cause of clutter in typical continental shelf environments [4, 5]. Based on the inverted parameters of fish such as neutral buoyancy depth, target strength and population density [6], scattered returns from fish distributions can be modeled and calibrated with measured returns. This model can then be extended to simulate wide-area images and movies, similar to those developed from collected data in the past [4, 5], showing temporal and spatial evolution of vast oceanic fish shoals.

We formulate a unified theory to model scattered returns from randomly distributed seafloor inhomogeneities in range-dependent ocean waveguides using the Rayleigh–Born approximation to

# Report Documentation Page

*Form Approved  
OMB No. 0704-0188*

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE <b>SEP 2011</b>	2. REPORT TYPE	3. DATES COVERED <b>00-00-2011 to 00-00-2011</b>			
4. TITLE AND SUBTITLE <b>Office of Naval Research Graduate Traineeship Award in Ocean Acoustics for Ankita Deepak Jain</b>		5a. CONTRACT NUMBER			
		5b. GRANT NUMBER			
		5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S)		5d. PROJECT NUMBER			
		5e. TASK NUMBER			
		5f. WORK UNIT NUMBER			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Massachusetts Institute of Technology, 77 Massachusetts Avenue, Room 5-212, Cambridge, MA, 02139</b>		8. PERFORMING ORGANIZATION REPORT NUMBER			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)			
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)			
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>4</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

Green's theorem [2]. This model will be implemented for a broadband source that uses matched-filter and beam forming to map the returns in range and bearing respectively. Calibration of modeled reverberation with data obtained during the OAWRS experiments enables estimation of seafloor parameters such as density, compressibility and coherence volume of inhomogeneities.

In the past, it has been shown that inexpensive underwater acoustic measurements can be used to determine the wind speed and classify the destructive power of a hurricane with greater accuracy than standard satellite remote sensing techniques [6]. The intensity of low frequency underwater sound measured directly below the hurricane is found to be approximately proportional to the cube of the local wind speed. In order to test this hypothesis, acoustic sensors will be deployed in the location on earth most frequented by hurricanes, and underwater noise data will be collected for a period of roughly one year. This noise data will then be correlated with local wind speed measurements and will be compared against data collected at another site, in the Mid Atlantic, to test the hypothesis developed by the PI.

## **WORK COMPLETED**

In order to characterize the spatial and temporal behavior of biological clutter, we model the scattered returns from schools of randomly distributed fish based on the inverted parameters of fish obtained from the data collected during the 2006 OAWRS experiment in the Gulf of Maine [6]. The total scattered field from a school is calculated using a full field matched filter model that is capable of charting the scattered returns in range. The modeled returns are evaluated at different stages of the shoal formation, i.e. before, during and after shoal formation is complete. These stages are marked by different fish population densities and distributions of fish in the water column as observed during the OAWRS experiments [4, 5]. The returns at these stages are found to be in agreement with those in observed data collected in the OAWRS experiment in the Gulf of Maine.

A full field matched filter approach is used to model scattered returns from random volume inhomogeneities in a range-dependent ocean environment [1]. Following the approach of Galinde et al. [2], a time harmonic approximation to the full field matched filter model is derived and is validated in Pekeris waveguide environment. The time-efficient time harmonic model is applied to the waveguide environments of the OAWRS experiments in the New Jersey continental shelf and the Gulf of Maine, and is used to estimate a combined seafloor parameter that is dependent on sediment properties such as density, compressibility and coherence volume of inhomogeneities. Reverberation data collected during the course of the two experiments for different source frequencies and bandwidths are compared with modeled returns to determine the spectral properties of this parameter. Since this parameter is independent of the propagation effect and is solely dependent on scattering properties of volume inhomogeneities in the seafloor, its estimation provides key information about reverberation in different waveguide environments.

To obtain more data relating underwater sound and wind power in hurricanes, acoustic hydrophones were deployed near Isla Socorro, Mexico as a part of a joint Ocean Acoustic Hurricane Classification experiment with the Mexican Navy. Isla Socorro, an island located a couple of hundred miles off the west coast of Mexico, is one of the most hurricane hit regions of the world and experiences an average of three hurricanes every year. In two experiments, one in 2007-08 and 2010-11, low frequency (<1000 Hz) underwater noise data was collected for a period of one year. Although, during both these hurricane seasons no hurricane passed near the island, we were still able to correlate the recorded underwater noise data with wind speeds, up to 15 m/s or 30 miles/hour, that were lower than those in

the hurricane range during the 2007-08 experiment. The noise intensity is found to be approximately proportional to the cube of the local wind speed [8], which is consistent with the results obtained by Wilson and Makris [7]. We found that low frequency underwater noise was often contaminated by anthropogenic airgun signals [8]. We find that during the periods when airgun contamination exists, it is still possible to extract information about natural geophysical noise by analyzing time series in between pulses, modeling airgun source transmission through the ocean, and correlating measured noise with local wind speed.

## IMPACT/APPLICATIONS

- A unified, range-dependent, broadband ocean reverberation model and calibration of modeled returns from seafloor sediments with past data will provide a tool for accurately estimating seafloor properties. This model helps distinguish between fluctuating returns due to clutter and statistically stationary seafloor reverberation.
- A model to calculate scattered returns from fish shoals will provide a tool to distinguish biological clutter from bathymetric features and intended targets. This model can also be used to study the spatio-temporal behavior of different fish species and marine organisms, which are a main cause of bioclutter in continental shelf environments.
- For wind speeds less than those in the hurricane range, low frequency underwater noise intensity is found to be approximately proportional to cube of the local wind speed. This result is consistent with the trend observed in the past for higher wind speeds in the hurricane range.

## REFERENCES

- [1] Anamaria Ignisca. *Analytic Model for Matched-filtered Scattered Intensity of volume inhomogeneities in an Ocean Waveguide Calibrated to Measured Seabed Reverberation*. Master of Science thesis, Massachusetts Institute of Technology, The Department of Mechanical Engineering, June 2011.
- [2] Ameya Galinde, Ninos Donabed, Mark Andrews, Sunwoong Lee, Nicholas C. Makris and Purnima Ratilal, "Range-dependent Waveguide Scattering Model Calibrated for Bottom Reverberation in Continental Shelf Environments" *J. Acoust. Soc. Am* 123, 1270-1281 (2008).
- [3] Mark Andrews, Zheng Gong, and Purnima Ratilal, "High-resolution population density imaging of random scatterers with the matched filtered scattered field variance", *J. Acoust. Soc. Am.* 126 1057-1068 (2009).
- [4] Nicholas C. Makris, Purnima Ratilal, Srinivasan Jagannathan, Zheng Gong, Mark Andrews, Ioannis Bertsatos, Olav Rune Godoe, Redwood W. Nero, J. Michael Jech, "Critical Population Density Triggers Rapid Formation of Vast Oceanic Fish Shoals", *Science*, Vol. 323, No. 5922, 1734-1737 (2009).
- [5] Nicholas C. Makris, Purnima Ratilal, Deanelle T. Symonds, Srinivasan Jagannathan, Sunwoong Lee and Redwood W. Nero, "Fish Population and Behavior Revealed by Instantaneous Continental Shelf-Scale Imaging," *Science*, 311, 660-663 (2006).

- [6] Zheng Gong, Mark Andrews, Srinivasan Jagannathan, Ruben Patel, J. Michael Jech, Nicholas C. Makris, Purnima Ratilal, "Low-frequency target strength and abundance of shoaling Atlantic herring *Clupea harengus* in the Gulf of Maine during the Ocean Acoustic Waveguide Remote Sensing (OAWRS) 2006 Experiment" *J. Acoust. Soc. Am.*, Vol. 127, 104-123 (2010).
- [7] Joshua D. Wilson and Nicholas C. Makris, "Quantifying hurricane destructive power, wind speed and air-sea material exchange with natural undersea sound," *Geophys. Res. Lett.* 35, L10603 (2008).
- [8] Ankita D. Jain, Srinivasan Jagannathan, Nicholas C. Makris Arturo C. Uribe , "Quantifying the contamination of natural low frequency ambient noise in the oceans by distant and pervasive anthropogenic airgun signals." *J. Acoust. Soc. Am.* (Cancun, Mexico, November 2010).